

IN THE CLAIMS

1. (Currently Amended) A medium for measuring the efficacy of a tumor therapy on single cell suspensions, comprising the essential amino acids, vitamins, salts and carbon donors, characterized in that the medium comprises from 0.1 to 1 mM buffer of pH 7.0 to 7.4, ~~5 to 20%~~ 5 to 20% by volume fetal calf serum, 4-6 g/l glucose and 2 to 5 mM glutamine as carbon source, wherein the medium does not contain carbon sources other than glucose and glutamine.
2. (Currently) The medium as claimed in claim 1, ~~characterized in that it further~~ comprises phosphate buffer as buffer.
3. (Canceled)
4. (Currently Amended) The medium as claimed in claim 1, ~~characterized in that~~ wherein the medium comprises 5 g/l glucose, 2 mM glutamine and 10% by volume fetal calf serum.
5. (Currently Amended) The medium as claimed in claim 2, ~~characterized in that~~ wherein the medium comprises 5 g/l glucose, 2 mM glutamine and 10% by volume fetal calf serum.
6. (Currently Amended) A method for measuring the efficacy of a tumor therapy on single cell suspensions of tumor cells by determining the acid formation in a medium in the presence and in the absence of a substance having cytostatic or cytotoxic activity, ~~characterized in that~~ wherein the measurement is carried out in a medium as claimed in claim 1.
7. (Currently Amended) The method as claimed in claim 6, ~~characterized in that~~ wherein the measurement is carried out by means of a pH electrode on which the single cell suspension is immobilized, with use of a flow cell through which the inventive medium flows.

8. (Currently Amended) The method as claimed in claim 6 or 7, ~~characterized in that~~
wherein the medium is pumped through the flow cell until a constant pH has been set up, and the
change in the pH with the medium stationary is then measured by measuring at intervals of 1.5 to
2.5 minutes, the medium is thereafter removed from the measurement cell, and the measurement
cycle is started from the beginning again until the pH change has been determined over a period
of 14 to 24 hours.

9. (Canceled)

10. (Currently Amended) The method as claimed in claim 9, ~~characterized in that~~
wherein the method is carried out in a multichannel instrument, with one channel being charged
by a medium without cytostatic and the other channels being charged with the medium
containing the cytostatic.